



National Health and Medical Research Council statement on electronic cigarettes: 2022 update

Becky Freeman¹ , Matthew J Peters^{2,3}, Renee Bittoun^{4,5}, Richard Brightwell⁶, Dallas R English⁷, David P Thomas⁸, Margaret FA Otlowski⁹, Nicholas A Zwar¹⁰ , Catherine Chamberlain^{11,12}

The National Health and Medical Research Council (NHMRC) of Australia is responsible for developing guidelines and evidence statements on a wide range of public and environmental health issues. Since 2015, this has included public health advice on electronic cigarettes (e-cigarettes). NHMRC issued a Chief Executive Officer (CEO) statement in 2017.¹ This article reports on the 2022 update to the NHMRC CEO statement on electronic cigarettes.

“E-cigarette” is an umbrella term for a range of devices that heat liquid to form an aerosol or vapour. The liquid in e-cigarettes (e-liquid) may contain nicotine and includes vegetable glycerine and propylene glycol.² Flavours and other chemicals are often added to the e-liquid. The resulting aerosol is quite different from cigarette smoke, with an absence or reduction of some smoke constituents but the addition of other chemicals and metals not associated with cigarette smoke. The device, the liquid mixture used, and variable heating temperatures can influence what is ultimately inhaled by e-cigarette users.³ E-cigarette devices have evolved from the early cigarette lookalike products to tank-based systems, where users add their own nicotine and liquids, to the current disposable and pre-filled pod products available.⁴

In Australia, the 2016 National Drug Strategy Household Survey (NDSHS)⁵ identified that 8.8% of the population aged over 14 years had “ever used” an e-cigarette.^{6,7} This increased to 11.3% of the total population aged over 14 years ever having used an e-cigarette in the 2019 NDSHS survey, and 6.9% of non-smokers over the age of 14 years reported ever having used an e-cigarette.^{7,8} Among those who had tried e-cigarettes, frequency of use also increased, with more people using them at least monthly (from 10.3% in 2016 to 17.9% in 2019).^{7,8} Adult daily e-cigarette use in Victoria increased from 0.9% to 2.4% from 2018–19 to 2022.⁹

Until early 2020, the majority of e-cigarette use in Australia was with modifiable tank-based systems. Since 2020, based on seized products and usage observations from a range of sources, disposable e-cigarettes now dominate.¹⁰ These are relatively low cost for first users, compared with tank systems; include nicotine in high concentration as nicotine salt; and come with colourful flavour descriptions designed to attract target users.¹¹ State-based data, such as in New South Wales, suggest that this change from tank-based to disposable high concentration devices has coincided with marked increase in e-cigarette use by young people.¹² Between 2020–21 and 2021–22, daily or occasional e-cigarette use in NSW by 16–24-year-olds increased from 11.1% to 16.5%.¹²

The combination of evolving product design, publication of new and emerging evidence, and growing concern among public

Abstract

Introduction: Electronic cigarette (e-cigarette) use in Australia has rapidly increased since the 2017 National Health and Medical Research Council (NHMRC) *Chief Executive Officer (CEO) statement on e-cigarettes*. The type of products available and the demographic characteristics of people using these products have changed. New evidence has been published and there is growing concern among public health professionals about the increased use, particularly among young people who do not currently smoke combustible cigarettes. The combination of these issues led NHMRC to review the current evidence and provide an updated statement on e-cigarettes. In this article, we describe the comprehensive process used to review the evidence and develop the 2022 NHMRC *CEO statement on electronic cigarettes*.

Main recommendations:

- E-cigarettes can be harmful; all e-cigarette users are exposed to chemicals and toxins that have the potential to cause adverse health effects.
- There are no health benefits of using e-cigarettes if you do not currently smoke tobacco cigarettes.
- Adolescents are more likely to try e-cigarettes if they are exposed to e-cigarettes on social media.
- Short term e-cigarette use may help some smokers to quit who have been previously unsuccessful with other smoking cessation aids. There are other proven safe and effective options available to help smokers to quit.

Changes in management as a result of this statement: The evidence base for the harms of e-cigarette use has strengthened since the previous NHMRC statement. Significant gaps in the evidence base remain, especially about the longer term health harms of using e-cigarettes and the toxicity of many chemicals in e-cigarettes inhaled as an aerosol.

health professionals about the increased use of e-cigarettes, particularly among young people who do not currently smoke combustible cigarettes, led NHMRC to review the current evidence and provide an updated NHMRC *CEO statement on electronic cigarettes* on their safety and health impacts.

As noted in the NHMRC Statement, nicotine is well understood to be the major addictive substance in tobacco cigarettes.¹³ Evidence for addictiveness of nicotine per se was therefore not under review. The NHMRC Statement also notes e-liquids can contain nicotine, even if labelled “nicotine-free”.¹⁴

The NHMRC Statement is intended to support the health of all Australians, not just current smokers. The Statement assists consumers and policy makers in understanding the current evidence relevant to the marketing and use of e-cigarettes, their impact on smoking initiation and cessation, uptake among youth, dual use with conventional tobacco products,

¹University of Sydney, Sydney, NSW. ²Macquarie University, Sydney, NSW. ³Concord Repatriation General Hospital, Sydney, NSW. ⁴Avondale University, Cooranbong, NSW. ⁵University of Notre Dame Australia, Sydney, NSW. ⁶Edith Cowan University, Perth, WA. ⁷Centre for MEGA Epidemiology, University of Melbourne, Melbourne, VIC. ⁸Menzies School of Health Research, Charles Darwin University, Darwin, NT. ⁹University of Tasmania, Hobart, TAS. ¹⁰Bond University, Gold Coast, QLD. ¹¹University of Melbourne, Melbourne, VIC. ¹²Judith Lumley Centre, La Trobe University, Melbourne, VIC. ✉ becky.freeman@sydney.edu.au • doi: [10.5694/mja2.52163](https://doi.org/10.5694/mja2.52163)

and their overall implications on individual and population health.


Methods

Overview of the revision process

The development of the 2022 NHMRC Statement took two years and followed best practice NHMRC guideline development

processes, as outlined in the 2016 NHMRC *standards for guidelines*¹⁵ and the NHMRC *Guidelines for guidelines handbook*.¹⁶ Key stages of the development process are outlined in Box 1. The process drew upon the diverse expertise of the NHMRC Electronic Cigarettes Working Committee¹⁷ (established at Step 1) to scope outcomes (Step 2), consider evidence reviews (Step 3), and the independent methodological review of those evidence reviews (Step 4). An evidence-to-decision process involved assessment of evidence quality and public health considerations using the

1 Overview of the development process of the 2022 National Health and Medical Research Council (NHMRC) statement on electronic cigarettes (e-cigarettes)



WHO-INTEGRATE framework¹⁸ (Step 5), targeted consultation (Step 6), and independent review (Step 7) before launch (Step 8).

The Committee considered a comprehensive mix of evidence when updating the NHMRC Statement. This included evidence statements and advice from other countries, such as the United States National Academies of Sciences, Engineering and Medicine¹⁹ and Surgeon General reports,^{20,21} and from Australia, such as the CSIRO (Commonwealth Scientific and Industrial Research Organisation) review²² and the National Industrial Chemicals Notification and Assessment Scheme review.²³ This was combined with evidence from commissioned evidence reviews (Step 3). In the evidence-to-decision process (Steps 5–7), relevant data on e-cigarette usage and sociodemographic and other features of the Australian community were also considered.

Populations and outcomes of interest (Step 2)

The Committee first established and determined which populations and outcomes to examine when updating the 2022 NHMRC Statement, in line with the intended use for broad public health. Populations considered were:

- never smokers (those who have never smoked a tobacco cigarette);
- former smokers (those who used to smoke tobacco cigarettes but have quit); and
- current smokers (those who currently smoke tobacco cigarettes).

Detailed outcomes were considered under three headings:

- e-cigarette use and tobacco smoking uptake and cessation;
- effects of e-cigarette advertising, promotion and sponsorship; and
- e-cigarette use and health outcomes.

The Committee did not examine or include evidence on the passive exposure to second hand aerosol from e-cigarettes or the appeal of e-cigarettes, including flavours, packaging design and price. The addictiveness of nicotine was not examined as this is well established and widely known.

Evidence reviews (Step 3)

In addition to considering existing international evidence, NHMRC and the Australian Government Department of Health and Aged Care also commissioned specific evidence reviews from external researchers. The Australian Government Department of Health and Aged Care commissioned the following evidence reviews:

- *E-cigarette use and combustible tobacco smoking uptake among non-smokers, including relapse of former smokers: umbrella review, systematic review and meta-analysis*²⁴;
- *Efficacy of e-cigarettes as aids to cessation of combustible tobacco smoking: updated evidence review*²⁵; and
- *Electronic cigarettes and health outcomes: systematic review of global evidence*.²⁶

In addition, NHMRC commissioned the following reviews:

- *Effects of e-cigarette advertising, promotion and sponsorship on people's attitudes, beliefs, perception, intentions and behaviours: a mixed methods systematic review*²⁷;

- *Supplementary report one: additional material on the review of evidence on the relationship of e-cigarette use to smoking behaviour, including uptake and cessation*²⁸; and
- *Supplementary report two: additional material on the review of evidence on the health outcomes of e-cigarette exposure*.²⁹

NHMRC also conducted the following scoping review to be included as part of the evidence base: *Inhalation toxicity of non-nicotine e-cigarette constituents: risk assessments, scoping review and evidence map* (toxicology report).³⁰

Methodological reviews (Step 4)

The research protocols for the reviews commissioned by NHMRC were reviewed by the Committee and independently methodologically reviewed before the reviews commenced. These protocols outlined the scope, research question and methodology.

All evidence reviews, supporting materials and supplementary reports produced were also independently methodologically reviewed to ensure that the methodological quality followed the systematic and rigorous approach documented in the review protocols. The reviews are publicly available from the NHMRC website.¹³

Evidence synthesis (Step 5)

The internationally recognised Grading of Recommendations, Assessment, Development and Evaluation (GRADE) methodology was used to assess certainty of the evidence.³¹ The GRADE methodology aims to improve transparency and consistency in reporting and decision making by assessing key aspects of the way studies are designed, run and analysed, which affects how certain (or confident) reviewers and committees can be that the results reported in studies are accurate. GRADE certainty ratings for the type of epidemiological evidence typical of broad public health exposures are generally low to very low. More highly rated study designs that are considered necessary for clinical practice guidelines, such as randomised controlled trials, are not appropriate for most public health exposures. This is chiefly because randomisation to many public health exposures of interest, such as smoking, alcohol, or early age at first pregnancy, is often not feasible or ethical. To acknowledge that well designed observational studies are often the best source of evidence on public health issues, Committee members agreed that bodies of evidence that consist of randomised controlled trials would assume an initial level of certainty of “high”, prospective cohort studies would assume an initial level of certainty of “moderate”, and all other observational studies would assume an initial level of certainty of “low”. The certainty of the evidence included in the evidence reviews varied across outcomes from moderate (for outcomes relating to the influence of the media on intention to use e-cigarettes in young people) to low or even very low for many of the other outcomes.

The available evidence had several limitations. E-cigarettes are relatively new products and the design and technology behind the devices and components used in e-cigarettes are constantly evolving. The wide variation of e-cigarettes, including manufacturing quality, setting customisation and variation in e-liquids used, makes it difficult to ascertain their safety and health impacts. Most of the direct evidence of safety and health impacts of e-cigarettes was limited to short term human epidemiological or clinical studies, with long term effects mostly unknown.

Most studies examining the toxicological constituents of e-liquids were *in vitro* or animal-based studies.³⁰ These study

designs have inherent limitations and are generally considered to provide less direct evidence when compared with human randomised controlled trials or long term epidemiological studies. However, *in vitro* or animal-based studies are useful when needing to understand the effects of an environmental exposure to a substance or when it is not ethical or possible to conduct human studies due to the toxic nature of the substances being examined or to control for confounding factors present in human populations.^{32,33}

The evidence on the impact of e-cigarette marketing on related behaviours, such as intention to use e-cigarettes or uptake/initiation, and on the impact of e-cigarette use on smoking uptake/initiation was mainly limited to international cross-sectional or cohort studies. Very few studies were conducted in Australia.²⁷ Contrastingly, most of the evidence examining the efficacy of e-cigarettes as a tobacco cigarette smoking cessation tool was from randomised controlled trials based on clinical settings with behavioural support, which may not be generalisable to real-world effectiveness of e-cigarettes in settings less supportive of cessation; these studies also had small sample sizes and short follow-up periods.²⁵

The WHO-INTEGRATE evidence-to-decision framework¹⁸ is designed for decision making at population levels and was used to develop the 2022 NHMRC Statement. The WHO-INTEGRATE framework was used to shape and write the 2022 NHMRC Statement, including what evidence was included and emphasised in the key messages.³⁴

Targeted consultation (Step 6) and independent expert review (Step 7)

Consultation is a core component of the NHMRC's guideline development process, contributing to accountability of the agency and independence of the advice. The draft 2022 NHMRC Statement and infographics underwent targeted consultation with federal, state and territory Chief Health/Medical Officers

over a two-week period in March 2022. Chief Health/Medical Officers were also asked to contribute Poisons Information Centre data on nicotine poisoning for inclusion in the Statement (if available) and to comment on two infographic concepts that summarised the Statement. Overall, Chief Health/Medical Officers were supportive of the Statement. In response to feedback, edits to the Statement were made, including the addition of a plain English summary.

After revisions in response to targeted consultation, the next version of the 2022 NHMRC Statement and associated resources underwent review by three independent experts to ensure the evidence on e-cigarettes was appropriately interpreted and synthesised. No errors of fact or significant omissions of evidence were identified. However, the phrasing of the 2022 NHMRC Statement and accompanying resources was clarified at several points. The final Statement and associated resources were released in June 2022 following consideration by the NHMRC Council.

Comparison with international guidance

The 2022 NHMRC Statement is broadly consistent with reports from a number of international health bodies published from 2020 onwards (Box 2). A significant factor contributing to variations between these statements and advice is the way that the questions have been framed,²² with some having a major focus on benefits associated with smoking cessation and others considering a broader range of issues including longer term harms and risks associated with use for purposes other than smoking cessation. The 2022 NHMRC Statement does not compare harms of e-cigarette use with smoking tobacco cigarettes, reflecting the purpose to support the health of all Australians and to avoid normalisation of nicotine. Relative harm from e-cigarette use compared with tobacco cigarettes is also difficult to quantify and varies depending on a number of factors such as conditions of use, e-cigarette device type and setting, e-liquid type and concentration, and frequency of e-cigarette and tobacco cigarette

2 Snapshot summary comparison of international guidance on electronic cigarettes (e-cigarettes)

Author	Guidance	Key messages
Belgian Superior Health Council	Report 9549: <i>Electronic cigarette: evolution</i> (2022) ³⁵	<ul style="list-style-type: none"> E-cigarettes are not recommended for non-smokers, especially young people. E-cigarettes are considered significantly less harmful than cigarettes. They are a better alternative to cigarettes for smokers and can be used as a smoking cessation aid.
World Health Organization	<i>Tobacco: e-cigarettes</i> (2022) ²	<ul style="list-style-type: none"> Both tobacco products and electronic nicotine devices (ENDS) pose risks to health. The safest approach is not to use either. To date, evidence on the use of ENDS as a cessation aid is inconclusive. In part due to the diversity of ENDS products and the low certainty surrounding many studies, the potential for ENDS to play a role as a population-level tobacco cessation intervention is unclear.
Scientific Committee on Health, Environmental and Emerging Risks (SCHEER)	<i>SCHEER scientific opinion on electronic cigarettes</i> (2021) ³⁶	<ul style="list-style-type: none"> There is moderate evidence that e-cigarettes are a gateway to smoking for young people and strong evidence that flavours have a relevant contribution for attractiveness of use of e-cigarettes and initiation. There is weak evidence for the support of e-cigarettes' effectiveness in helping smokers to quit and the evidence on smoking reduction is assessed as weak to moderate.
United States Preventive Services Task Force	<i>Interventions for tobacco smoking cessation in adults, including pregnant persons</i> (2021) ³⁷	<ul style="list-style-type: none"> The current evidence is insufficient to assess the balance of benefits and harms of e-cigarettes for tobacco cessation in adults, including pregnant persons.
Irish Health Research Board	<i>Electronic cigarette use and tobacco cigarette smoking initiation in adolescents: an evidence review</i> (2020) ³⁸	<ul style="list-style-type: none"> This review found a fourfold association between ever using e-cigarettes and initiating smoking tobacco cigarettes in adolescents.
	<i>Electronic cigarette and smoking cessation: an evidence review</i> (2020) ³⁹	<ul style="list-style-type: none"> The systematic review and network meta-analysis of electronic nicotine delivery systems (e-cigarettes) versus therapies usually given for smoking cessation showed that there is no evidence of a difference in effect on incidences of smoking cessation. There is a low level of certainty in these results due to low successful event rates and high rates lost to follow-up in all studies.

use.^{3,40} The absolute risks of e-cigarettes cannot be determined at the present time and longer term effects, of importance for young people who become dependent, remain unclear.¹⁹ As such, the 2022 NHMRC Statement can be characterised as supporting a continuation of the precautionary approach.

Key messages

Following this extensive evidence synthesis, targeted consultation and expert review, a precautionary approach to e-cigarettes, particularly for people who have never smoked, remains appropriate. The key messages in the 2022 NHMRC Statement¹³ include:

- E-cigarettes can be harmful; all e-cigarette users are exposed to chemicals and toxins that have the potential to cause adverse health effects.
- E-cigarette-related poisonings have substantially increased over the past five years. E-cigarette-related calls to Australian Poisons Information Centres have more than doubled between 2020 and 2021.
- There are no health benefits of using e-cigarettes if you do not currently smoke tobacco cigarettes.
- People who have never smoked may be more likely to take up tobacco smoking if they use e-cigarettes.
- Adolescents are more likely to try e-cigarettes if they are exposed to e-cigarettes on social media.
- Short term e-cigarette use may benefit smokers if they are able to quit smoking and have been previously unsuccessful with other smoking cessation aids.
- There are other proven safe and effective options available to help smokers quit.

Following the development and publication of the NHMRC Statement, a media release and associated social media content (Box 3) promoting and explaining the Statement were issued. The 2022 NHMRC Statement is an essential synthesis of the current evidence that can now be used by both state and federal health departments in guiding e-cigarette regulations. Further research is needed in many areas, including on the longer term impacts of e-cigarette use on the health of young people.

The NHMRC 2022 *CEO statement on electronic cigarettes* and supporting literature are available at www.nhmrc.gov.au/ecigs.

3 Summary infographic of the National Health and Medical Research Council 2022 *CEO statement on electronic cigarettes (e-cigarettes)*



Source: Figure reproduced with permission from the National Health and Medical Research Council.¹³ ♦

Acknowledgements: We would like to show our gratitude and pay our respects to our colleague, Dr Kerry Nugent, who died on 29 December 2022 after a brief illness. Dr Nugent was a highly valued member of the Electronic Cigarettes Working Committee. We are especially grateful for his contributions to the development of the National Health and Medical Research Council (NHMRC)'s 2022 *CEO statement on electronic cigarettes*, including his sound advice and knowledge on the toxicological aspects of e-liquids and e-cigarettes. We also acknowledge the contribution of Jennifer Savenake, Director (Public Health), and the NHMRC project team throughout the development of the NHMRC 2022 *CEO statement on electronic cigarettes*.

Open access: Open access publishing facilitated by The University of Sydney, as part of the Wiley - The University of Sydney agreement via the Council of Australian University Librarians.

Competing interests: The authors were reimbursed for their time to attend NHMRC Electronic Cigarettes Working Committee meetings. All committee member declarations can be found in full here: <https://www.nhmrc.gov.au/about-us/leadership-and-governance/committees/electronic-cigarettes-working-committee>.

Provenance: Not commissioned; externally peer reviewed. ■

© 2023 The Authors. *Medical Journal of Australia* published by John Wiley & Sons Australia, Ltd on behalf of AMPCo Pty Ltd.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

1 National Health and Medical Research Council. *CEO statement on electronic cigarettes (e-cigarettes)*. Canberra: NHMRC, 2017. <https://www.nhmrc.gov.au/sites/default/files/documents/attachments/statement-electronic-cigarettes.pdf> (Viewed Oct 2023).

2 World Health Organization. *Tobacco: e-cigarettes*. Geneva: WHO, 2022. [https://www.who.int/news-room/questions-and-answers/item/tobacco-e-cigarettes#:~:text=Electronic%20cigarettes%20\(or%20e%20cigarettes,of%20nicotine%20and%20harmful%20emissions](https://www.who.int/news-room/questions-and-answers/item/tobacco-e-cigarettes#:~:text=Electronic%20cigarettes%20(or%20e%20cigarettes,of%20nicotine%20and%20harmful%20emissions) (viewed Feb 2023).

3 World Health Organization. *Electronic nicotine delivery systems and electronic non-nicotine delivery systems (ENDS/ENNDs)*. Geneva: WHO, 2016. [https://www.who.int/publications/m/item/electronic-nicotine-delivery-systems-and-electronic-non-nicotine-delivery-systems-\(ends-ennds\)](https://www.who.int/publications/m/item/electronic-nicotine-delivery-systems-and-electronic-non-nicotine-delivery-systems-(ends-ennds)) (viewed Feb 2023).

4 Centers for Disease Control and Prevention. *About electronic cigarettes (e-cigarettes)*. Atlanta (GA): Office on Smoking and Health, 2023. https://www.cdc.gov/tobacco/basic_infor

2016. [https://www.who.int/publications/m/item/electronic-nicotine-delivery-systems-and-electronic-non-nicotine-delivery-systems-\(ends-ennds\)](https://www.who.int/publications/m/item/electronic-nicotine-delivery-systems-and-electronic-non-nicotine-delivery-systems-(ends-ennds)) (viewed Feb 2023).

4 Centers for Disease Control and Prevention. *About electronic cigarettes (e-cigarettes)*. Atlanta (GA): Office on Smoking and Health, 2023. https://www.cdc.gov/tobacco/basic_infor

- mation/e-cigarettes/about-e-cigarettes.html (viewed Feb 2023).
- 5 Australian Institute of Health and Welfare. National Drug Strategy Household Survey, 2016 [supplementary data tables: chapter 3, Tobacco; table 3.16: Lifetime use of electronic cigarettes (e-cigarettes), by age and smoker status, people aged 12 years or older, 2013 and 2016 (per cent)]. Canberra: AIHW, 2016. <https://www.aihw.gov.au/getmedia/c679f184-ca65-4fc6-9a00-dbb4feb56c02/chapter-3-tobacco.xlsx.aspx> (viewed Oct 2023).
 - 6 Tobacco in Australia. Prevalence of e-cigarette use [table 18.3.1]. Melbourne: Cancer Council, 2019. https://www.tobaccoinaustralia.org.au/chapter-18-e-cigarettes/18-3-extent#_ENREF_5 (viewed Oct 2023).
 - 7 Australian Institute of Health and Welfare. National Drug Strategy Household Survey 2019 [Cat. No. PHE 270]. Canberra: AIHW, 2020. <https://www.aihw.gov.au/reports/illicit-use-of-drugs/national-drug-strategy-household-survey-2019/contents/summary> (viewed Feb 2023).
 - 8 Australian Institute of Health and Welfare. National Drug Strategy Household Survey, 2019 [data tables: National Drug Strategy Household Survey 2019: 2 Tobacco smoking, supplementary tables. Table 2.19: Lifetime use of electronic cigarettes (e-cigarettes), by age and smoker status, 2016 to 2019 (per cent)]. Canberra: AIHW, 2020. <https://www.aihw.gov.au/getmedia/e83fc585-87e9-466b-8f63-6821a74b5528/aihw-phe-260-2-Tobacco-smoking-tables.xlsx.aspx> (viewed Oct 2023).
 - 9 Bayly M, Mitsopoulos E, Durkin S, Scollo M. E-cigarette use and purchasing behaviour among Victorian adults: findings from the 2018–19 and 2022 Victorian Smoking and Health Surveys [prepared for Quit Victoria]. Melbourne: Centre for Behavioural Research in Cancer, Cancer Council Victoria; 2022. https://www.cancer.vic.org.au/downloads/cbrc/R22_MB_E-cigarette%20use%20and%20purchasing%20behaviour%20among%20Victorian%20adults.pdf (viewed Oct 2023).
 - 10 Watts C, Egger S, Dessaix A, et al. Vaping product access and use among 14–17-year-olds in New South Wales: a cross-sectional study. *Aust N Z J Public Health* 2022; 46: 814–820.
 - 11 NSW Health. The facts about vaping. Sydney: NSW Government, 2022 <https://www.health.nsw.gov.au/tobacco/Factsheets/vaping-teachers-types.pdf> (viewed Feb 2023).
 - 12 HealthStats NSW. E-cigarette use by electronic cigarette use and age (years) [NSW Population Health Survey (SAPHaRI), Centre for Epidemiology and Evidence, NSW Ministry of Health]. Sydney: NSW Health, 2022 [https://www.healthstats.nsw.gov.au/#/indicator?name=-beh-smo-ecig-phs&location=NSW&view=Trend&measure=prevalence&groups=Age%20\(years\),Electronic%20cigarette%20use&compare=Electronic%20cigarette%20use,Age%20\(years\)&filter=Electronic%20cigarette%20use,Current%20user,Ever%20used&filter=Age%20\(years\),All%20ages,16-24%20years](https://www.healthstats.nsw.gov.au/#/indicator?name=-beh-smo-ecig-phs&location=NSW&view=Trend&measure=prevalence&groups=Age%20(years),Electronic%20cigarette%20use&compare=Electronic%20cigarette%20use,Age%20(years)&filter=Electronic%20cigarette%20use,Current%20user,Ever%20used&filter=Age%20(years),All%20ages,16-24%20years) (viewed Feb 2023).
 - 13 National Health and Medical Research Council. CEO statement on electronic cigarettes. Canberra: NHMRC, 2022. <https://www.nhmrc.gov.au/health-advice/all-topics/electronic-cigarettes/ceo-statement> (viewed Feb 2023).
 - 14 Chivers E, Janka M, Franklin P, et al. Nicotine and other potentially harmful compounds in “nicotine-free” e-cigarette liquids in Australia. *Med J Aust* 2019; 210: 127–128. <https://www.mja.com.au/journal/2019/210/3/nicotine-and-other-potentially-harmful-compounds-nicotine-free-e-cigarette>
 - 15 National Health and Medical Research Council. 2016 NHMRC standards for guidelines. Canberra: NHMRC, 2016. <https://www.nhmrc.gov.au/guidelinesforguidelines/standards> (viewed Feb 2023).
 - 16 National Health and Medical Research Council. Guidelines for guidelines handbook. Canberra: NHMRC, 2016 <https://www.nhmrc.gov.au/guidelinesforguidelines> (viewed Feb 2023).
 - 17 National Health and Medical Research Council. Electronic Cigarettes Working Committee. Canberra: NHMRC, 2022 <https://www.nhmrc.gov.au/about-us/leadership-and-governance/committees/electronic-cigarettes-working-committee> (viewed Feb 2023).
 - 18 Rehfuess EA, Stratil JM, Scheel IB, et al. The WHO–INTEGRATE evidence to decision framework version 1.0: integrating WHO norms and values and a complexity perspective. *BMJ Global Health* 2019; 4 (Suppl): e000844.
 - 19 National Academies of Sciences, Engineering and Medicine. Public health consequences of e-cigarettes. Washington, DC: National Academy Press, 2018. <https://nap.nationalacademies.org/catalog/24952/public-health-consequences-of-e-cigarettes> (viewed Sept 2023).
 - 20 US Department of Health and Human Services. E-cigarette use among youth and young adults: a report of the Surgeon General. Atlanta (GA): US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2016. https://www.cdc.gov/tobacco/data_statistics/sgr/e-cigarettes/pdfs/2016_sgr_entire_report_508.pdf (viewed Feb 2023).
 - 21 US Department of Health and Human Services. Smoking cessation: a report of the Surgeon General. Rockville (MD): US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2020. <https://www.hhs.gov/sites/default/files/2020-cessation-sgr-full-report.pdf> (viewed Feb 2023).
 - 22 Byrne S, Brindal E, Williams G, et al. E-cigarettes, smoking and health: a literature review update. *CSIRO*, 2018. <https://www.csiro.au/-/media/BF/Files/E-cigarettes/E-cigarettes-Consolidated-Final-Report240618-pdf.pdf> (viewed Sept 2023).
 - 23 Australian Government Department of Health, National Industrial Chemicals Notification and Assessment Scheme. Non-nicotine liquids for e-cigarette devices in Australia: chemistry and health concerns. Canberra: Commonwealth of Australia, 2019. <https://www.industrialchemicals.gov.au/sites/default/files/2020-08/Non-nicotine%20liquids%20for%20e-cigarette%20devices%20in%20Australia%20chemistry%20and%20health%20concerns%20%5BPDF%201.21%20MB%5D.pdf> (viewed Feb 2023).
 - 24 Baenziger ON, Ford L, Yazidjoglou A, Joshy G, Banks E. E-cigarette use and combustible tobacco cigarette smoking uptake among non-smokers, including relapse in former smokers: umbrella review, systematic review and meta-analysis. *BMJ Open* 2021; 11: e045603.
 - 25 Yazidjoglou A, Ford L, Baenziger O, et al. Efficacy of e-cigarettes as aids to cessation of combustible tobacco smoking: updated evidence review — final report prepared for the Australian Government Department of Health. Canberra: National Centre for Epidemiology and Population Health, Australian National University; 2021. <https://openresearch-repository.anu.edu.au/handle/1885/247864> (viewed Feb 2023).
 - 26 Banks E, Yazidjoglou A, Brown S, et al. Electronic cigarettes and health outcomes: systematic review of global evidence — report to the Australian Department of Health. Canberra: National Centre for Epidemiology and Population Health, Australian National University; 2022. https://openresearch-repository.anu.edu.au/bitstream/1885/262914/1/Electronic%20cigarettes%20health%20outcomes%20review_2022_WCAG.pdf (viewed Feb 2023).
 - 27 Moola S, Tyagi J, Miller M, et al. Effects of e-cigarette advertising, promotion, and sponsorship on people’s attitudes, beliefs, perceptions, intentions, and behaviours: a mixed methods systematic review — evidence evaluation report. Sydney: George Institute for Global Health, 2021. <https://www.nhmrc.gov.au/file/18976/download?token=PMTRLJaL> (viewed Feb 2023).
 - 28 Banks E, Beckwith K, Yazidjoglou A, et al. Supplementary report one: additional material on the review of evidence on the relationship of e-cigarette use to smoking behaviour, including uptake and cessation — final report prepared for the National Health and Medical Research Council. Canberra: National Centre for Epidemiology and Population Health, Australian National University; 2021. https://www.nhmrc.gov.au/file/18290/download?token=_CnGFuna (viewed Feb 2023).
 - 29 Banks E, Yazidjoglou A, Beckwith K, et al. Supplementary report two: additional material on the review of evidence on the health outcomes of e-cigarette exposure. Canberra: National Centre for Epidemiology and Population Health, Australian National University; 2021. <https://www.nhmrc.gov.au/file/18288/download?token=ILAY7j1> (viewed Feb 2023).
 - 30 National Health and Medical Research Council. Inhalation toxicity of non-nicotine e-cigarette constituents: risk assessments, scoping review and evidence map. Canberra: NHMRC, 2022. https://www.nhmrc.gov.au/sites/default/files/documents/attachments/ecigarettes/Scoping_review_on_the_inhalation_toxicity_of_non-nicotine_e-cigarette_constituents.pdf (viewed Feb 2023).
 - 31 Schünemann H, Brożek J, Guyatt G, Oxman A. Handbook for grading the quality of evidence and the strength of recommendations using the GRADE approach. GRADE Working Group, 2013. <https://gdt.gradepro.org/app/handbook/handbook.html#h.9rdbelsnu4iy> (viewed Sept 2023).
 - 32 Woodruff TJ, Sutton P. Navigation Guide Work Group. An evidence-based medicine methodology to bridge the gap between clinical and environmental health sciences. *Health Aff (Millwood)* 2011; 30: 931–937.
 - 33 Woodruff TJ, Sutton P. The Navigation Guide systematic review methodology: a rigorous and transparent method for translating environmental health science into better health outcomes. *Environ Health Perspect* 2014; 122: 1007–1014.
 - 34 Santesso N, Glenton C, Dahm P, et al. GRADE guidelines 26: informative statements to communicate the findings of systematic reviews of interventions. *J Clin Epidemiol* 2020; 119: 126–135.
 - 35 Conseil Supérieur de la Santé. [Electronic cigarette: evolution] [French]. Brussels: CSS, 2022. <https://www.health.belgium.be/sites/>

default/files/uploads/fields/fpshealth_theme_file/20220616_css-9549_avis_e-cigarette_vweb_0.pdf (viewed Sept 2023).

36 Scientific Committee on Health, Environmental and Emerging Risks (SCHEER). SCHEER scientific opinion on electronic cigarettes. Brussels: European Commission, 2021. https://health.ec.europa.eu/system/files/2022-08/scheer_o_017.pdf (viewed Sept 2023).

37 US Preventive Services Task Force; Krist AH, Davidson KW, Mangione CM, et al. Interventions for tobacco smoking cessation in adults,

including pregnant persons: US Preventive Services Task Force recommendation statement. *JAMA* 2021; 325: 265-279.

38 O'Brien D, Long J, Lee C, et al. Electronic cigarette use and tobacco cigarette smoking initiation in adolescents: an evidence review. Dublin: Health Research Board, 2020. https://www.hrb.ie/fileadmin/2._Plugin_related_files/Publications/2020_publication-related_files/2020_HIE/Evidence_Centre/Electronic_cigarette_use_and_tobacco_cigarette_smoking_initiation_in_adolescents.pdf (viewed Sept 2023).

39 Quigley J, Kennelly H, Lee C, et al. Electronic cigarettes and smoking cessation: an evidence review. Dublin: Health Research Board, 2020. https://www.hrb.ie/fileadmin/2._Plugin_related_files/Publications/2020_publication-related_files/2020_HIE/Evidence_Centre/Electronic_cigarettes_and_smoking_cessation_systematic_evidence_review.pdf (viewed Sept 2023).

40 Lindson-Hawley N, Hartmann-Boyce J, Fanshawe TR, et al. Interventions to reduce harm from continued tobacco use. *Cochrane Database Syst Rev* 2016; (10): CD005231. ■