One remarkable feature of the present system is its capacity for activating both molecular nitrogen and hydrogen under mild conditions. The incorporation of hydrogen into the titanium complex as a hydride and the succeeding transfer to magnesium without reaction with the reduced nitrogen is noteworthy in view of the susceptibility of the nitrogen–titanium complexes to attack by protic reagents.

A similar system composed of VCl_3 and magnesium in THF is also active in nitrogen fixation.

(*Received, April 28th,* 1969; *Com.* 570.)

[1] This system was reported by S. Go, K. Soga, Y. Hattori, and T. Keii at the 21st Annual Meeting of the Chemical Society of Japan, Osaka, Japan, April 1968.
[2] M. E. Vol'pin, M. A. Ilatovskaya, L. V. Kosyakova, and V. B. Shur, *Chem. Comm.*, 1968, 1074.
[3] Recently the formation of sodium hydride by a titanium compound–sodium naphthalenide–H_2 system was reported; E. E. van Tamelen and R. B. Fechter, *J. Amer. Chem. Soc.*, 1968, **90**, 6854.

The Application of Simple Hückel Theory to Metal Trisacetylacetonates

By E. A. MAGNUSSON and K. A. THOMSON
(*Avondale College, Cooranbong, New South Wales, Australia,* 2265)
and A. G. WEDD*†
(*Department of Chemistry, University of Tasmania, Hobart, Tasmania, Australia*)

Summary Previous Hückel calculations applied to metal trisacetylacetonates are shown to be in error and are repeated and extended in the light of recent ionisation potential data, indicating that the approach is quantitatively inadequate for the present system.

SIMPLE Hückel theory has been employed by Barnum[1] to estimate the effect of metal–ligand (M–L) π-bonding on the electronic spectra[2] of a series of tervalent trisacetylacetonates, $M(acac)_3$ (M = Ti, V, Cr, Mn, Fe, Co). Coulomb and exchange parameters were estimated from electronegativity and bond energy data, respectively, with the exception of the exchange parameter, β_{MO}, for the metal–oxygen π-interaction, which was left as a variable parameter. The original paper[1] should be consulted for background, method and notation. The treatment resulted in remarkably good correlations with the experimental band positions.

Important recent developments have been mass spectral investigations[4,5] of the trisacetylacetonates, in which the first appearance potentials of the $M(acac)_3^+$ ions are identified with the ionisation potentials of $M(acac)_3$ and compared with those predicted by the Barnum calculation and extensions[5] of the latter to Hacac, $Al(acac)_3$ and complexes of closely related ligands. The calculations fail to reproduce even the qualitative trends of the mass spectral data (which are similar to those for the uncomplexed ligands) and it is suggested[5] that Koopmans' theorem may not apply to this system.

In the original calculation,[1] the eigenvalues for the functions of E symmetry were obtained by diagonalising the 6×6 energy matrix formed from the single given set of E-type symmetry orbitals. Unfortunately, however, this set of orbitals, ϕ, is not orthogonal to the two further sets, ϕ' and ϕ'', which are easily set down by cyclic permutation of basis functions. Thus, while correct doubly-degenerate eigenvalues might be obtained by diagonalising the 12×12 matrix formed from two of these linearly dependent sets, those obtained in the original paper are incorrect as terms off-diagonal with respect to the sets are omitted. The most straightforward step[6] to obtain proper orthogonal functions is to consider the two sets ϕ and $(\phi' - \phi'')$.

Suitable combinations of these can be taken, resulting in two new sets, one of which is symmetric, the other antisymmetric with respect to one of the C_2 axes. This transformation factorises the 12×12 matrix and one of the three possible sets of un-normalised orbitals is given in the Table.

TABLE

E-type symmetry orbitals for $M(acac)_3$[a]

Symmetric w.r.t. C_2

$$2d_{xy} - d_{xz} - d_{yz}$$
$$2\phi_A - 2\phi_E - \phi_F + \phi_J - \phi_K + \phi_O$$
$$2\phi_B - 2\phi_D - \phi_G + \phi_I - \phi_L + \phi_N$$
$$\phi_H - \phi_M$$
$$\phi_G + \phi_I - \phi_L - \phi_N$$
$$\phi_F + \phi_J - \phi_K - \phi_O$$

Antisymmetric w.r.t. C_2

$$d_{yz} - d_{xz}$$
$$2\phi_A + 2\phi_E - \phi_F - \phi_J - \phi_K - \phi_O$$
$$2\phi_B + 2\phi_D - \phi_G - \phi_I - \phi_L - \phi_N$$
$$2\phi_C - \phi_H - \phi_M$$
$$\phi_F - \phi_J - \phi_K + \phi_O$$
$$\phi_G - \phi_I - \phi_L + \phi_N$$

[a] For details of the notation, see ref. 1.

While the simple Hückel theory is much too approximate to be useful in interpreting detailed electronic spectra, it was plausible that the recent mass spectral data[4,5] might be more amenable to correlation with a simple one-electron energy level scheme and therefore the calculations were repeated. The values employed by Barnum for the parameters α_C, α_O, β_{CO} and β_{CC} were again used. As Forster's Hückel calculation on the acetylacetonate ion indicated that the results were sensitive to parameter choice, the values recommended by Streitwieser[8] (and used by Forster) were also used, and side-by-side comparisons of results using identical β_{MO} and α_M values made. Barnum's differential electronegativity correction formulae were followed. The problem was extended to two dimensions by variation of α_M, as well as β_{MO}. Barnum estimated his α_M values from neutral atom electronegativities and cited the Pauling electroneutrality principle as justification. It is now recognised[11,12] that this principle may not be obeyed as closely as previously thought. The valence state ionisation energy for the *d*-orbital of the free atom ground-state configuration of the individual metals[13] was used as an initial estimate of α_M, which was then systematically varied over a range of 30 kK. This range covered Barnum's values, but no ligand-field splitting corrections were included. The exchange parameter β_{MO} was again left as an adjustable parameter. The calculation procedure requires only the diagonalisation of a matrix.

For most values of α_M in the above range, a value of β_{MO} could be found which would fit the first appearance potentials of the mass spectra of the complexes to the highest occupied MO.‡ However, no consistent trend in the values of the parameters was apparent. There appears to be little advantage in tabulating specific examples of the calculations.

Despite the failure, one important conclusion of the original analysis[1] can be corroborated. Thus the interaction of the metal A_1 orbital with the lower, filled, bonding and higher, empty, antibonding ligand A_1 orbitals leaves its energy largely unchanged. This suppression of the effect of π-interaction (assuming it is significant) would show why the position of the acetylacetonate ion in the spectrochemical series indicates little or no π-antibonding effects.

While the simple MO picture may provide a reasonable pictorial description of the relative order of the occupied levels of highest energy in the ground states of these complexes,[3] the gross approximations inherent in the Hückel approach, plus the apparent non-applicability of Koopmans' theorem,[5] do not allow a more quantitative description of the properties considered here.

(*Received, May 5th,* 1969; *Com.* 628.)

† Present address: Division of Pure Chemistry, National Research Council, Ottawa, Canada.

‡ Using octahedral group notation for convenience, the energies of the e_g orbitals were estimated (ref. 2) as energy less negative by a value Δ than those of the t_{2g} orbitals.

[1] D. W. Barnum, *J. Inorg. Nuclear. Chem.*, 1961, **22**, 183.
[2] D. W. Barnum, *J. Inorg. Nucl. Chem.*, 1961, **21**, 221.
[3] J. P. Fackler, jun., *Progr. Inorg. Chem.*, 1966, **7**, 361.
[4] G. M. Bancroft, C. Reichert, and J. B. Westmore, *Inorg. Chem.*, 1968, **7**, 870.
[5] S. M. Schildcrout, R. G. Pearson, and F. E. Stafford, *J. Amer. Chem. Soc.*, 1968, **90**, 15, 4006.
[6] F. A. Cotton, "Chemical Applications of Group Theory," Interscience, 1963.
[7] L. S. Forster, *J. Amer. Chem. Soc.*, 1964, **86**, 3001.
[8] A. Streitwieser, jun., "Molecular Orbital Theory for Organic Chemists," Wiley, New York, 1961.
[9] C. K. Jørgensen, *Acta Chem. Scand.*, 1962, **16**, 2406.
[10] C. K. Jørgensen, "Absorption Spectra and Chemical Bonding in Complexes," Pergamon, Oxford, 1962.
[11] C. K. Jørgensen, "Orbitals in Atoms and Molecules," Academic Press, London, 1962.
[12] C. K. Jørgensen, S. M. Horner, W. Hatfield, and S. Y. Tyree, jun., *Internat. J. Quantum Chem.*, 1967, **1**, 191.
[13] H. Basch, A. Viste, and H. B. Gray, *Theor. Chim. Acta*, 1965, **3**, 458.

Trigonal Prismatic Co-ordination in the *cis,cis*-1,3,5-Tris(pyridine-2-carboxaldimino)cyclohexanezinc(II) Ion

By W. O. GILLUM, J. C. HUFFMAN, W. E. STREIB,* and R. A. D. WENTWORTH
(*Department of Chemistry, Indiana University, Bloomington, Indiana* 47401)

Summary A single-crystal *X*-ray diffraction study has shown the Zn^{II} complex of *cis,cis*-1,3,5-tris(pyridine-2-carboxaldimino)cyclohexane to have trigonal prismatic co-ordination.

HERETOFORE, authenticated examples of trigonal prismatic co-ordination were limited to compounds which exclusively contain sulphur-bonded metal atoms from the second and third transition series.[1,2] Our interests in hindered ligand systems[3] has led to a three-dimensional *X*-ray study of the Zn^{II} complex of *cis,cis*-1,3,5-tris(pyridine-2-carboxaldimino)cyclohexane, and we find, as shown in Figure 1, that trigonal prismatic co-ordination can occur with nitrogen-bonded metal atoms of the first transition series as well.

The anhydrous perchlorate crystallizes in the space group $P2_1/c$ with $a = 13{\cdot}440 \pm 0{\cdot}008$, $b = 13{\cdot}738 \pm 0{\cdot}006$, $c = 15{\cdot}279 \pm 0{\cdot}008$ Å, $\beta = 109°\ 53' \pm 1'$, $Z = 4$, $D_m = 1{\cdot}67$ g.cm.$^{-3}$, and $D_c = 1{\cdot}65$ g.cm.$^{-3}$. A fully automatic Picker diffractometer, equipped with a highly oriented graphite monochromator, was used to collect a 2-fold redundant set of data totalling 11,369 reflections using the standard θ–2θ scan technique and Mo-K_α radiation. The set of 2185 unique reflections with $I_{obs} > 2{\cdot}33\sigma_I$, (where σ_I is the standard error in I_{obs} based on counting statistics)



FIGURE 1. *Structure of the* cis,cis-1,3,5-*tris*(*pyridine-2-carboxaldimino*)*cyclohexanezinc*(II) *ion. The pseudo-threefold axis is horizontal and in the plane of the paper.*