

11-2015

# Risk of Organism Acquisition From Prior Room Occupants: A Systematic Review and Meta-Analysis

Brett G. Mitchell

Avondale College of Higher Education, [brett.mitchell@avondale.edu.au](mailto:brett.mitchell@avondale.edu.au)

Stephanie Dancer

National Health Service Scotland, [Stephanie.Dancer@lanarkshire.scot.nhs.uk](mailto:Stephanie.Dancer@lanarkshire.scot.nhs.uk)

Malcolm Anderson

Avondale College, [malcolm.anderson@avondale.edu.au](mailto:malcolm.anderson@avondale.edu.au)

Emily Dehn

Avondale College of Higher Education, [s11176409@student.avondale.edu.au](mailto:s11176409@student.avondale.edu.au)

Follow this and additional works at: [https://research.avondale.edu.au/nh\\_conferences](https://research.avondale.edu.au/nh_conferences)



Part of the [Nursing Commons](#)

---

## Recommended Citation

Mitchell, B. G., Dancer, S., Anderson, M., & Dehn, E. (2015, November). *Risk of organism acquisition from prior room occupants: A systematic review and meta-analysis*. Poster presented at International Australasian College of Infection Prevention and Control (ACIPC) Conference, Hobart, Australia.

This Conference Proceeding is brought to you for free and open access by the Faculty of Nursing and Health at ResearchOnline@Avondale. It has been accepted for inclusion in Nursing and Health Conference Papers by an authorized administrator of ResearchOnline@Avondale. For more information, please contact [alicia.starr@avondale.edu.au](mailto:alicia.starr@avondale.edu.au).

# Risk of organism acquisition from prior room occupants: A systematic review and meta-analysis

A/Professor Brett Mitchell <sup>1-2</sup>

Dr Stephanie Dancer <sup>3</sup>

Dr Malcolm Anderson <sup>1</sup>

Emily Dehn <sup>1</sup>

<sup>1</sup> Avondale College; <sup>2</sup> Australian Catholic University;

<sup>3</sup> Hairmyres Hospital, East Kilbride

# Disclosures

- Brett Mitchell Chair of Scientific Committee
- Brett Mitchell Interim Editor-in-Chief of Infection Disease and Health
- Study funded via an Avondale scholarship

# Background

- Environment plays a role in facilitating the transmission of important pathogens
- Organisms survive
- Studies have shown that if a patient is admitted to a room where the prior occupant was colonised or infected with a hospital pathogen, there is an increased risk of the next patient acquiring the same organism

# Purpose of systematic review

- Determine whether being admitted to a room where the prior occupant was colonized or infected with an organism increases the risk of acquiring that organism.
  - Explore differences in the risk of acquisition between Gram-positive and Gram-negative organisms.

# Methods: Search strategy

- Systematic review and meta-analysis
- PROSPERO: CRD42015016273
- Medline/PubMed, Cochrane and CINHAL
- Observational studies, last 30 years
- Must have examined exposure or acquisition in a hospitalized population where the prior room occupant was colonized or infected with a specific organism

# Methods

## Organisms

- *Acinetobacter*
- *Escherichia coli*
- *Klebsiella*
- *Pseudomonas*
- *Enterobacter*
- *Citrobacter*
- *Proteus*
- *Serratia*
- *Enterococcus*
- *C.difficile*
- *S.aureus* & VRE


## Exclusions

- Conference abstracts,
- Letters to editors
- Reviews
- Papers written in languages other than English

# Methods

- Assessed risk of bias (ROB) and quality using modified version of NOS (Wells et al., 2014)
- Random effects model used for meta-analysis
- Heterogeneity assessed using  $I^2$  statistic






# Results

Study (lead author)	Year	Study duration	Study setting (country)	Study design	Organisms
Huang	2005	20 months	USA	Cohort	VRE, MRSA
Mitchell	2014	24 months	Australia	Cohort	MRSA
Datta	2011	20 months	USA	Cohort	VRE, MRSA
Ajao	2013	93 months	USA	Cohort	ESBL-producing Gram negative
Drees	2008	14 months	USA	Cohort	VRE
Nseir	2011	12 months	France	Cohort	<i>A.baumannii</i> , ESBL-producing Gram negative, <i>P. aeruginosa</i>
Shaughnessy	2011	16 months	USA	Cohort	<i>C. difficile</i>

# Results

- 4,643 'exposed' patients → 287 (6.2% )  
acquired the same species of organism.
- 34,886 'unexposed' patients → 1,112 (3.2%)
- Pooled acquisition OR for all the organisms  
included in the six studies was 2.14 (95% CI =  
1.65–2.77)
- Pooled acquisition OR for Gram-negative  
organisms was 2.65(95% CI = 2.02–3.47) and  
1.89 (95% CI = 1.62–2.21) for Gram-positive  
organisms

Study or Subgroup	Decreased acquisition		Control		Weight	Odds Ratio M-H, Random, 95% CI
	Events	Total	Events	Total		
Huang (MRSA)	57	1454	248	8697	16.2%	1.39 [1.04, 1.86]
Nseir (ESBL producing Gram Neg)	8	50	50	461	0.0%	1.57 [0.70, 3.52]
Huang (VRE)	58	1291	256	9058	16.2%	1.62 [1.21, 2.16]
Ajao (Klebsiella sp. or Escherichia coli)	32	648	235	8723	14.2%	1.88 [1.29, 2.74]
Nseir (Pseudomonas)	21	85	61	426	10.4%	1.96 [1.12, 3.45]
Drees (VRE)	19	138	31	500	9.7%	2.42 [1.32, 4.43]
Shaughnessy (Clostridium difficile)	10	91	77	1679	8.3%	2.57 [1.28, 5.15]
Mitchell (MRSA)	74	884	163	5344	16.4%	2.90 [2.18, 3.86]
Nseir (Acinetobacter)	16	52	41	459	8.6%	4.53 [2.32, 8.86]
<b>Total (95% CI)</b>		<b>4643</b>		<b>34886</b>	<b>100.0%</b>	<b>2.14 [1.65, 2.77]</b>
Total events	287		1112			
Heterogeneity: Tau <sup>2</sup> = 0.09; Chi <sup>2</sup> = 21.32, df = 7 (P = 0.003); I <sup>2</sup> = 67%						
Test for overall effect: Z = 5.74 (P < 0.00001)						



# Results - sub analysis

- Gram negative organisms, *A. baumannii* had the highest odds ratio (OR 4.53 = 2.32-8.86).
- Further sub-analyses → no differences:
  - *C.difficile* against the MRSA studies;
  - MRSA against the VRE studies;
  - *Klebsiella* species and *E.coli* ESBL-producing Gram-negative bacilli with *Pseudomonas aeruginosa* against *Acinetobacter baumannii*.
  - In acquisition between ESBL producing organisms and MRSA or VRE.

# Key points

- Admission to a room previously occupied by a patient infected and/or colonised with a specific pathogen is a risk factor for acquisition.
- Regardless of the organism (species) the risk of acquisition increases
- Greater pooled acquisition rate for Gram-negative organisms

# Implications

- ICPs - understanding and managing the risks associated with the determination of room placement.
- Knowing the status of the prior room occupant may serve as important information in decision-making
- Current cleaning practices fail to reduce the risk of acquisition.
- Supports the need to improve hospital design
- Wider public → our study opens up a discussion about what is deemed acceptable risk.

# Limitations

- Constrained by the limitations of the individual studies reviewed
  - inability to conduct meta-regression
  - different approaches to testing the efforts of the participants, potential variations in microbiological testing methods
  - the presumption of acquisition based on epidemiological evidence
  - the inability to account for colonisation pressure



# Conclusion

- Prior room occupancy is a risk factor for acquisition
- Renewed interest and emphasis on hospital cleaning, and particularly discharge or terminal cleaning.

(Journal of Hospital Infection, 91(3):211-7)

# References

- Ajao AO, Johnson K, Harris AD, et al. Risk of acquiring extended spectrum b-lactamase-producing Klebsiella species and Escherichia coli from prior room occupants in the intensive care unit. Infect Control Hosp Epidemiol 2013;34:453e458.
- Datta R, Platt R, Yokoe DS, Huang SS. Environmental cleaning intervention and risk of acquiring multidrug-resistant organisms from prior room occupants. Archs Intern Med 2011;171:491e494.
- Drees M, Snyderman DR, Schmid CH, et al. Prior environmental contamination increases the risk of acquisition of vancomycin-resistant enterococci. Clin Infect Dis 2008;46:678e685.
- Mitchell BG, Digney W, Ferguson JK. Prior room occupancy increases risk of methicillin-resistant Staphylococcus aureus acquisition. Healthcare Infect 2014;19:135e140.
- Nseir S, Blazejewski C, Lubret R, Wallet F, Courcol R, Durocher A. Risk of acquiring multidrug-resistant Gram-negative bacilli from prior room occupants in the intensive care unit. Clin Microbiol Infect 2011;17:1201e1208.
- Shaughnessy MK, Micielli RL, DePestel DD, et al. Evaluation of hospital room assignment and acquisition of Clostridium difficile infection. Infect Control Hosp Epidemiol 2011;32:201e206.
- Wells G, Shea B, O'Connell D, et al. The NewcastleOttawa Scale (NOS) for assessing the quality of nonrandomised studies in metaanalyses 2014. Available at: [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp)